


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

win 55212-2 cyclodextrin concentration

Search

[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)
**Scholar** All articles - **Recent articles** Results 11 - 20 of about 34 for win 55212-2 cyclodextrin concentration with Sefsearch on. (0.19 seconds)

Topical WIN55212-2 Alleviates Intraocular Hypertension in Rats Through a CB1 Receptor Mediated ... - all 3 versions »

MH Oltmanns, SS Samudre, IG Castillo, A Hosseini, ... - Journal of Ocular Pharmacology and Therapeutics, 2008 - liebertonline.com

 ... From 60–120 min, a twentyfold in- crease in **WIN 55212-2** was seen in the urine. **WIN**
**55212-2 concentration** in the blood was below de- tectable limits. ...

[Related Articles](#) - [Web Search](#) - [BL Direct](#)
A search for presynaptic 3-adrenoceptors in the rat

D •laszczyk, A Zakrzewska, G Kwolok, B Malinowska, ... - Fundamental &amp; Clinical Pharmacology, 2005 - ingentaconnect.com

 ... a 19% w/v solution of **cyclodextrin (WIN 55212-2)** or in a mixture of saline ... experiments)

 or saline (in situ experiments) to obtain the **concentration** required. ...

[Related Articles](#) - [Web Search](#)
Effects of Cannabinoids on Sympathetic and Parasympathetic Neuroeffector Transmission in the Rabbit ... - all 4 versions »

B Szabo, U Nordheim, N Niederhoffer - Journal of Pharmacology and Experimental Therapeutics, 2001 - ASPET

 ... for cannabinoids, 2-hydroxypropyl- -**cyclodextrin**, was injected ... and the plasma noradrenaline **concentration** after systemic ... receptor agonist, **WIN 55212-2**, and is ...

[Cited by 23](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)
Role of CGRP and GABA in the hypotensive effect of intrathecally administered anandamide to ... - all 4 versions »

M del Carmen Garcia, E Adler-Graschinsky, S Maris ... - European Journal of Pharmacology, 2006 - Elsevier

 ... The final **concentration** of ethanol for the two drugs was 14% (v/v). **WIN 55212-2**

 was dissolved in 10% w/v solution of hydroxypropyl- $\beta$ -**cyclodextrin**. ...

[Cited by 1](#) - [Related Articles](#) - [Web Search](#)
R (+)-methanandamide-induced cyclooxygenase-2 expression in H4 human neuroglioma cells: possible ... - all 4 versions »

B Hinz, R Ramer, K Eichele, U Weinzierl, K Brune - Biochemical and Biophysical Research Communications, 2004 - Elsevier

 ... the influence of methyl- $\beta$ -**cyclodextrin** (MCD), a ... was associated with a **concentration**-dependent inhibition ... C. Giaume, Anandamide and **WIN 55212-2** inhibit cyclic ...

[Cited by 8](#) - [Related Articles](#) - [Web Search](#)
Constrained analogs of CB-1 antagonists: 1, 5, 6, 7-Tetrahydro-4H-pyrrolo [3, 2-c] pyridine-4-one ... - all 3 versions »

RA Smith, Z Fathi, SE Brown, S Choi, J Fan, S ... - Bioorganic &amp; Medicinal Chemistry Letters, 2007 - Elsevier

... 2. Effect of the pyrrolopidridone hydrazide 44 on food intake in overnight-fasted

Wistar rats, after dosing at 10 mg/kg po as a suspension in 30% **cyclodextrin** ...

[Cited by 2](#) - [Related Articles](#) - [Web Search](#)

**Cannabinoids for Therapeutic Use: Designing Systems to Increase Efficacy and Reliability.** - [all 2 versions »](#)

F Grotenhermen - American Journal of Drug Delivery, 2004 - drugdelivery.adisonline.com

... The area under the plasma **concentration**-time curve (AUC ... 69,70) the synthetic derivative

**WIN 55212-2**, [71] and ... applied through the use of **cyclodextrin** technology. ...

[Related Articles](#) - [Web Search](#)

**Role of cortical cannabinoid CB1 receptor in conditioned taste aversion memory** - [all 6 versions »](#)

T Kobilio, S Hazvi, Y Dudai - European Journal of Neuroscience, 2007 - Blackwell Synergy

... **2-hydroxypropyl-1-β-cyclodextrin** (Cavasol®, Sigma ... with saline to a final **concentration** of 0.9 ... FA & Takahashi, RN (2006) **WIN 55212-2** impairs contextual ...

[Cited by 4](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

**Optimization of imidazole amide derivatives as cannabinoid-1 receptor antagonists for the treatment ...** - [all 3 versions »](#)

RA Smith, Z Fathi, F Achebe, C Akuche, SE Brown, S ... - Bioorganic & Medicinal Chemistry Letters, 2007 - Elsevier

... unfortunately all of these compounds were found to provide quite poor plasma exposure from oral dosing (10 mg/kg po, typically in 30% **cyclodextrin** suspension). ...

[Web Search](#)


**Cannabinoids: reward, dependence, and underlying neurochemical mechanisms—a review of recent ...** - [all 3 versions »](#)

G Tanda, SR Goldberg - Psychopharmacology, 2003 - Springer

... Veh THC vehicle (**b-cyclodextrin**) administration ... by a progressive decrease of cocaine **concentration** in the ... CB1 receptor agonist **WIN 55,212-2** showed reinforcing ...

[Cited by 85](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

Key authors: [A Gifford](#) - [S Gatley](#) - [A Makriyannis](#) - [N Volkow](#) - [N Niederhoffer](#)



Result Page: [Previous](#) [1](#) [2](#) [3](#) [4](#) [Next](#)

win 55212-2 cyclodextrin concentrat

Search

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

